




Ain Shams University

Ain Shams Engineering Journal

www.elsevier.com/locate/asej
www.sciencedirect.com



ENGINEERING PHYSICS AND MATHEMATICS

Solving generalized quintic complex Ginzburg–Landau equation by homotopy analysis method

Soheila Naghshband ^{a,b}, Mohammad Ali Fariborzi Araghi ^{a,*}

^a Department of Mathematics, Central Tehran Branch, Islamic Azad University, Tehran, Iran

^b Department of Mathematics, West Tehran Branch, Islamic Azad University, Tehran, Iran

Received 19 June 2015; revised 18 December 2015; accepted 28 January 2016

KEYWORDS

Generalized quintic complex Ginzburg–Landau (GCGL) equation;
 Homotopy analysis method (HAM);
 Nonlinearity;
 h -curve;
 Variational iteration method (VIM)

Abstract In this paper, the generalized quintic complex Ginzburg–Landau equation is considered to be solved, by means of the homotopy analysis method (HAM). Two examples are solved to illustrate the efficiency of the proposed method. By plotting the h -curve of the examples, the region of convergence is determined.

© 2016 Faculty of Engineering, Ain Shams University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The cubic–quintic complex Ginzburg–Landau equation is a continuous approximation to the dynamics of the field in a passively mode-locked laser. It has also been proven to be useful in describing important phenomena such as ultrashort pulse propagation in optical transmission lines with spectral filtering and erbium-doped fiber amplifiers, the GCGL equation was used successfully in modeling of other nonequilibrium processes in physics. The quintic terms in the equation describe important physics, which is lacking in other models in the

literature [3]. It arises in a variety of settings, including nonlinear optics, fluid dynamics, chemical physics, mathematical biology, condensed matter physics, and statistical mechanics. In fluid dynamics the GCGL is found, for example, in the study of Poiseuille flow, the nonlinear growth of convection rolls in the Rayleigh–Benard problem and Taylor–Couette flow. In this case, the bifurcation parameter R plays the role of a Reynolds number. The equation also arises in the study of chemical systems governed by reaction–diffusion equations. The GCGL equation plays the role of a simplified set of fluid dynamic equations.

The Ginzburg–Landau equation has been used to study many practical problems such as chemical turbulence, Poiseuille flow, Taylor–Couette flow, Rayleigh–Benard convection, reaction–diffusion systems, nonlinear optics, and hydrodynamical stability problems. It exhibits rich dynamics and has become a paradigm for the transition to spatio-temporal chaos. Also some other details about the other different types of Ginzburg equation can be considered in [3,7,8,10,24,25,26,27,31,33,34].

* Corresponding author. Tel.: +98 2188385773; fax: +98 2188385777.

E-mail addresses: m_fariborzi@iauctb.ac.ir, fariborzi.araghi@gmail.com (M.A. Fariborzi Araghi).

☆ Peer review under responsibility of Ain Shams University.



Production and hosting by Elsevier

<http://dx.doi.org/10.1016/j.asej.2016.01.015>


2090-4479 © 2016 Faculty of Engineering, Ain Shams University. Production and hosting by Elsevier B.V.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Please cite this article in press as: Naghshband S, Fariborzi Araghi MA, Solving generalized quintic complex Ginzburg–Landau equation by homotopy analysis method, Ain Shams Eng J (2016), <http://dx.doi.org/10.1016/j.asej.2016.01.015>

Table 1 The results of example 1 via the HAM at $x = 1$.

t	0	0.2	0.4	0.6	0.8	1
Error	0	1.1102e-016	1.1102e-016	1.4983e-016	1.4895e-015	4.7744e-014

**Fig. 1** The h -curve of $w_{tt}(0.5, 1)$ given by 7-approximation of the HAM.

We consider generalized quintic complex Ginzburg–Landau equation with the form

$$\frac{\partial w}{\partial t} = (1 + ia)\frac{\partial^2 w}{\partial x^2} + Rw - (1 + ib)|w|^{2n}w - (1 + id)|w|^{4n}w, \quad n \geq 1, \\ w(x, 0) = f(x), \quad i^2 = -1, \quad (1)$$

where a, b, d, R are real constants and $w = w(x, t)$ is a complex unknown function and t is a nonnegative real value, x is a real value [3,9,13,14,29,33,34]. Wazwaz studied Eq. (1) by using the separation of variables method in [33].

In this work, the HAM is being considered to obtain the approximate solution of Eq. (1). This method is a powerful analytical method to solve the nonlinear problems and was first introduced and used by Liao [23]. Recently, this method has been well applied to solve many types of problems in science and engineering [1,2,4–6,11,12,17–19,28,30,32]. Homotopy analysis method contains an auxiliary parameter h which provides a simple way to adjust and control the convergent region and the rate of convergence of the series solution. HAM bears a very rapid convergence of the solution series in most cases, usually only a few iterations with satisfactory approximate solution, as well [20–23]. Total description of this paper is as follows: In Section 2, some preliminaries are given,

and in Section 3, the main idea of this paper is brought, and finally in Section 4, two examples are solved and h -curves are plotted to show the region of convergence.

2. Preliminaries

Let the following differential equation:

$$N[w(x, t)] = 0,$$

where N is a nonlinear operator, x and t denote the independent variables and w is an unknown function. Via the HAM, the zeroth-order deformation equation is:

$$(1 - q)L[\Phi(x, t, q) - w_0(x, t)] = qhH(x, t)N[\Phi(x, t, q)], \quad (2)$$

where $q \in [0, 1]$ is the embedding parameter, $h \neq 0$ is an auxiliary parameter, L is an auxiliary linear operator and $H(x, t)$ is an auxiliary function. $\Phi(x, t, q)$ is an unknown function and $w_0(x, t)$ is an initial guess of $w(x, t)$. It is clear, if $q = 0$ and $q = 1$ then:

$$\Phi(x, t, 0) = w_0(x, t), \quad \Phi(x, t, 1) = w(x, t),$$

respectively. Therefore, when q increases from 0 to 1, the solution $\Phi(x, t, q)$ varies from $w_0(x, t)$ to the exact solution $w(x, t)$. By Taylor's theorem, it can be expanded $\Phi(x, t, q)$ in a power series of the embedding parameter q as comes:

$$\Phi(x, t, q) = w_0(x, t) + \sum_{m=1}^{\infty} w_m(x, t)q^m, \quad (3)$$

where

$$w_m(x, t) = \frac{1}{m!} \left. \frac{\partial^m \Phi(x, t, q)}{\partial q^m} \right|_{q=0}. \quad (4)$$

Let the initial guess $w_0(x, t)$, the auxiliary linear operator L , the nonzero auxiliary parameter h and the auxiliary function $H(x, t)$ be properly chosen so that the power series Eq. (3) converges at $q = 1$, then, it can be seen:

$$w(x, t) = w_0(x, t) + \sum_{m=1}^{\infty} w_m(x, t), \quad (5)$$

which must be the solution of the original nonlinear equation. Now, we define the following set of vectors:

$$\vec{w}_n = \{w_0(x, t), w_1(x, t), \dots, w_n(x, t)\}. \quad (6)$$

Table 2 The errors of the example 1 via HAM at (2, 0.7).

n	Approximation at (2, 0.7)	by HAM ($h = -1$)	Error
4	-9.028807597814920e-001	+ 4.281032621672038e-001i	1.393798067426522e-003
8	-9.040720445762246e-001	+ 4.27379933574138e-001i	1.109812228348067e-007
12	-9.040721420156820e-001	+ 4.273798802345464e-001i	1.554304121895804e-012
16	-9.040721420170611e-001	+ 4.273798802338299e-001i	1.570092458683775e-016




Fig. 2 Imaginary and real parts of exact and approximation solutions of Example 1 when $x = -1$ and $m = 7$.




Fig. 3 The h -curve for $w_t(1, 0)$ given by 6-approximation ($m = 6$).

By differentiating the zeroth order deformation Eq. (2) m times with respect to the embedding parameter q and then setting $q = 0$ and finally dividing by $m!$, we will have the following m th order deformation equation:

$$L[w_m(x, t) - \chi_m w_{m-1}(x, t)] = hH(x, t)R_m(\vec{w}_{m-1}), \quad (7)$$

where

$$R_m(\vec{w}_{m-1}) = \frac{1}{(m-1)!} \left. \frac{\partial^{m-1} N[\Phi(x, t, q)]}{\partial q^{m-1}} \right|_{q=0}, \quad (8)$$

and

$$\chi_m = \begin{cases} 0 & m \leq 1, \\ 1 & m > 1. \end{cases} \quad (9)$$

It should be mentioned that $w_m(x, t)$ for $m \geq 1$ is governed by the linear Eq. (7) with linear boundary conditions that come from the original problem. For more details about the HAM, we refer the reader to [20,21].

3. Main idea

We consider Eq. (1) as follows:

$$\begin{aligned} \frac{\partial w}{\partial t} &= (1 + ia) \frac{\partial^2 w}{\partial x^2} + Rw - (1 + ib)w^{n+1}\bar{w}^n \\ &\quad - (1 + id)w^{2n+1}\bar{w}^{2n}n \geq 1, \end{aligned} \quad (10)$$

$$w(x, 0) = f(x), \quad i^2 = -1,$$

and

$$L[\Phi(x, t, q)] = \frac{\partial \Phi(x, t, q)}{\partial t}, \quad L(c) = 0, \quad (11)$$

where c is a real constant,

$$\begin{aligned} N[\Phi(x, t, q)] &= \frac{\partial \Phi(x, t, q)}{\partial t} - (1 + ia) \frac{\partial^2 \Phi(x, t, q)}{\partial x^2} \\ &\quad - R\Phi(x, t, q) + (1 + ib)\Phi^{n+1}(x, t, q)\bar{\Phi}^n(x, t, q) + (1 + id)\Phi^{2n+1}(x, t, q)\bar{\Phi}^{2n}(x, t, q), \end{aligned} \quad (12)$$

and $H(x, t) = 1$. The zeroth-order deformation equation is:

$$(1 - q)L[\Phi(x, t, q) - w_0] = qhN[\Phi(x, t, q)]. \quad (13)$$

Also, the m th-order deformation equation:

$$L[w_m - \chi_m w_{m-1}] = hR_m(\vec{w}_{m-1}), \quad (14)$$

where




Fig. 4 Imaginary parts of the approximate and the exact(left) solutions of Example 2 when $m = 6$.




Fig. 5 Real parts of the approximate and the exact(left) solutions of Example 2 when $m = 6$.

$$\begin{aligned}
 R_m(\vec{w}_{m-1}) &= \frac{\partial w_{m-1}}{\partial t} - (1+ia) \frac{\partial^2 w_{m-1}}{\partial x^2} - R w_{m-1} + (1+ib) \sum_{k_1=0}^{m-1} \sum_{k_2=0}^{k_1} \sum_{k_3=0}^{k_1-k_2} \cdots \sum_{k_{n+1}=0}^{k_1-k_2-\cdots-k_n} \sum_{\alpha_1=0}^{m-1-k_1} \sum_{\alpha_2=0}^{m-1-k_1-\alpha_1} \cdots \sum_{\alpha_{n-1}=0}^{m-1-k_1-\alpha_1-\cdots-\alpha_{n-2}} w_{k_2}, \\
 w_{k_3} \cdots w_{k_{n+1}} w_{(k_1-k_2-\cdots-k_{n+1})} \bar{w}_{\alpha_1} \bar{w}_{\alpha_2} \cdots \bar{w}_{\alpha_{n-1}} \bar{w}_{(m-1-k_1-\alpha_1-\cdots-\alpha_{n-1})} &+ (1+id) \sum_{k_1=0}^{m-1} \sum_{k_2=0}^{k_1} \sum_{k_3=0}^{k_1-k_2} \cdots \sum_{k_{2n+1}=0}^{k_1-k_2-\cdots-k_{2n}} \sum_{\alpha_1=0}^{m-1-k_1} \sum_{\alpha_2=0}^{m-1-k_1-\alpha_1} \cdots \sum_{\alpha_{2n-1}=0}^{(m-1-k_1-\alpha_1-\cdots-\alpha_{2n-2})} w_{k_2}, \\
 w_{k_3} \cdots w_{k_{2n+1}} w_{(k_1-k_2-\cdots-k_{2n+1})} \bar{w}_{\alpha_1} \bar{w}_{\alpha_2} \cdots \bar{w}_{\alpha_{2n-1}} \bar{w}_{(m-1-k_1-\alpha_1-\cdots-\alpha_{2n-1})}. &
 \end{aligned} \tag{15}$$

So,

$$w_m = \chi_m w_{m-1} + h \int_0^t R_m(\vec{w}_{m-1}) dt + c, \quad m \geq 1. \quad (16)$$

Now, in order to solve Eq. (10), with other semi-analytical method, we consider the variational iteration method (VIM) [15,16] to compare with the HAM. Let the following nonlinear differential equation is given:

$$Lw(x, t) + Nw(x, t) = g(x, t),$$

where L and N are linear and nonlinear operators respectively, and g is inhomogeneous term. The correction functional can be written as follows:

$$w_{m+1}(x, t) = w_m(x, t) + \int_0^t \lambda(\xi) (Lw_m(\xi) + N\tilde{w}_m(\xi) - g(\xi)) d\xi, \\ m \geq 0,$$

where λ is general Lagrange multiplier which can be calculated by variational theory, that it means $\delta \tilde{w}_m = 0$, and the exact solution is determined by $w = \lim_{m \rightarrow \infty} w_m$ [15,33]. By applying the VIM to solve Eq. (10), the correction functional of Eq. (10) can be considered as follows:

$$w_{m+1} = w_m + \int_0^t \lambda(\xi) \left(\frac{\partial w_m}{\partial \xi}(x, \xi) - (1 + ia) \frac{\partial^2 \tilde{w}_m}{\partial x^2}(x, \xi) - R\tilde{w}_m(x, \xi) + (1 + ib) \tilde{w}_m^{n+1} \tilde{w}_m^n(x, \xi) + (1 + id) \tilde{w}_m^{2n+1} \tilde{w}_m^{2n}(x, \xi) \right) d\xi. \quad (17)$$

To make this correction functional stationary, by applying, $\delta w_m(x, 0) = 0$, we obtain:

$$\delta w_{m+1}(x, t) = \delta w_m(x, t) + \int_0^t \lambda(\xi) \left(\frac{\partial}{\partial \xi} \delta w_m(x, \xi) \right) d\xi = 0.$$

So, the stationary conditions can be written as follows:

$$\lambda'(\xi) = 0, \quad 1 + \lambda(\xi)|_{\xi=t} = 0.$$

Thus, the Lagrange multiplier can be obtained as $\lambda = -1$, also and the following iteration formula is as follows:

$$w_{m+1} = w_m - \int_0^t \left(\frac{\partial w_m}{\partial \xi}(x, \xi) - (1 + ia) \frac{\partial^2 w_m}{\partial x^2}(x, \xi) - R w_m(x, \xi) + (1 + ib) w_m^{n+1} \tilde{w}_m^n(x, \xi) + (1 + id) w_m^{2n+1} \tilde{w}_m^{2n}(x, \xi) \right) d\xi. \quad (18)$$

4. Sample examples

In this section, two generalized quintic complex Ginzburg–Landau equations are solved via the HAM based on Eqs. (15) and (16) and the results are shown in the tables. Also, the h -curves of each example are plotted. The programs and figures have been provided by Matlab package.

Example 1. Consider the following PDE:

$$w_t = (1 + i)w_{xx} + 3w - (1 + 2i)|w|^2 w - (1 - 4i)|w|^4 w, \quad w(x, 0) = e^{ix}, \quad i^2 = -1.$$

By the HAM, we get

$$w_0(x, t) = e^{ix}, \\ w_1(x, t) = -hte^{xi}i, \\ w_2(x, t) = -(hte^{xi}(2hi + ht + 2i))/2, \\ w_3(x, t) = -(hte^{xi}(-h^2 t^2 i + 6h^2 t + 6h^2 i + 6ht + 12hi + 6i))/6, \dots$$

setting $h = -1$, we obtain

$$w_0(x, t) = e^{ix}, \\ w_1(x, t) = ite^{xi}, \\ w_2(x, t) = -(t^2 e^{xi})/2 = \frac{(it)^2 e^{ix}}{2!}, \\ w_3(x, t) = -(t^3 e^{xi}i)/6 = \frac{(it)^3 e^{ix}}{3!}, \dots$$

so, we have

$$w(x, t) = w_0(x, t) + w_1(x, t) + w_2(x, t) + \dots \\ = e^{ix} + \frac{(it)^2 e^{ix}}{2!} + \frac{(it)^3 e^{ix}}{3!} + \dots \\ = e^{ix} \left(1 + \frac{(it)^2}{2!} + \frac{(it)^3}{3!} + \dots \right) = e^{i(x+t)},$$

which is the exact solution of this example.

Table 1 shows the errors of the HAM at $x = 1$ when $m = 15$ and $h = -1$, for different values of t . Also, Fig. 1 shows the h -curve of w_{tt} obtained from the 7-approximation via the proposed method.

Table 2 shows the convergence of the HAM at the point (2, 0.7) and $m = 4, 8, 12, 16$, also the error column is calculated by $|\sum_{i=0}^m w_i - w|$. Fig. 2 shows the Imaginary and real parts of exact and approximation solutions of the proposed method, when $x = -1$ and different values of t , for Example 1.

Applying Eq. (18) as the variational iteration method, the following results are obtained:

$$w_0(x, t) = e^{ix}, \quad w_1(x, t) = e^{ix} + ite^{xi}, \quad w_2(x, t) = e^{ix} + ite^{xi} - (t^2 e^{xi})/2, \dots$$

so, it can be seen:

$$\lim_{m \rightarrow \infty} w_m = e^{ix} \left(1 + \frac{(it)^2}{2!} + \frac{(it)^3}{3!} + \dots \right) = e^{i(x+t)}.$$

It means that, the VIM is also able to obtain the solution of the equation by a convergent recursive scheme and this result can be found via the HAM in the special case of $h = -1$.

Example 2. Consider the following PDE:

$$w_t = (1 + 3i)w_{xx} + 3w - (1 + i)|w|^4 w - (1 - 6i)|w|^8 w, \quad w(x, 0) = e^{-ix}, \quad i^2 = -1,$$

applying the HAM, we get

$$w_0(x, t) = e^{-ix}, \\ w_1(x, t) = -2hte^{-xi}i, \\ w_2(x, t) = -2hte^{-xi}(hi + ht + i), \\ w_3(x, t) = -(4hte^{-xi})(-h^2 t^2 i + 3h^2 t + (3h^2 i)/2 + 3ht + 3hi + (3i)/2))/3, \dots,$$

setting $h = -1$, we obtain

$$w_0(x, t) = e^{-ix},$$

$$w_1(x, t) = 2ite^{-xi},$$

$$w_2(x, t) = -2t^2e^{-xi} = \frac{(2it)^2e^{-ix}}{2!},$$

$$w_3(x, t) = -4(t^3e^{-xi})/3 = \frac{(2it)^3e^{-ix}}{3!}, \dots,$$

so, we have

$$\begin{aligned} w(x, t) &= w_0(x, t) + w_1(x, t) + w_2(x, t) + \dots \\ &= e^{-ix} + \frac{(2it)^2e^{-ix}}{2!} + \frac{(2it)^3e^{-ix}}{3!} + \dots \\ &= e^{-ix} \left(1 + \frac{(2it)^2}{2!} + \frac{(2it)^3}{3!} + \dots \right) = e^{i(2t-x)}, \end{aligned}$$

which is the exact solution of this example.

Fig. 3 presents the h -curve of for $w_t(1, 0)$. Also, Figs. 4 and 5 compare the imaginary and real parts of the approximate and the exact solution, where $-2 \leq x \leq 2$.

5. Conclusion

In this paper, the homotopy analysis method was used to solve the generalized quintic complex Ginzburg–Landau equation. In order to show the importance and applicability of the proposed method, two examples by plotting of the h -curves and some related numerical results were thoroughly illustrated. Also, in an example, it can be observed that the VIM is also a proper and efficiency method to solve Eq. (10), whereas it is not able to control and adjust the region of convergence. Consequently, the HAM can be applied to solve the generalized quintic GCGL equation as a reliable and valid scheme.

Acknowledgements

The authors would like to thank the reviewers of this paper for their valuable recommendations and also the Islamic Azad University, central Tehran branch, for their support during this research.

References

- [1] Abbasbandy S. The application of the homotopy analysis method to nonlinear equations arising in heat transfer. *Phys Lett A* 2006;360(1):109–13.
- [2] Abbasbandy S. The application of homotopy analysis method to solve a generalized Hirota–Satsuma coupled KdV equation. *Phys Lett A* 2007;361(6):478–83.
- [3] Akhmediev NN, Afanasjev VV. Stability of the pulselike solutions of the quintic complex Ginzburg–Landau equation. *J Opt Soc Am B* 1996;13(7):1439–49.
- [4] Alizadeh-Pahlavan A, Aliakbar V, Vakili-Farahani F, Sadeghy K. MHD flows of UCM fluids above porous stretching sheets using two-auxiliary-parameter homotopy analysis method. *Commun Nonlinear Sci Numer Simul* 2009;14(2):473–88.
- [5] Alomari AK, Noorani MSM, Nazar R. Explicit series solution of some linear and non-linear Schrodinger equations via the homotopy analysis method. *Commun Nonlinear Sci Numer Simul* 2009;14:1196–207.
- [6] Alomari AK, Noorani MSM, Nazar R. Comparison between the homotopy analysis method and homotopy perturbation method to solve coupled Schrodinger-KdV equation. *J Appl Math Comput* 2009;31(1–2):1–12.
- [7] Aranson IS, Kramer L. The world of the complex Ginzburg–Landau equation, 2001.
- [8] Biazar J, Partovi M, Ayati Z. Approximating solutions for Ginzburg–Landau equation by HPM and ADM. *Appl Appl Math* 2010;1672–81.
- [9] Charbonneau J. Introduction to the Ginzburg–Landau equations. Vancouver (BC, Canada, V6T 1Z1): Department of Physics and Astronomy, University of British Columbia; 2005 [November 22].
- [10] Eguiluz V, Hernandez-Garcia E, Piro O. Boundary effects in the complex Ginzburg–Landau equation. *Internat J Bifur Chaos* 1999;9(11):2209–14.
- [11] Fariborzi Araghi MA, Fallahzadeh A. On the convergence of the homotopy analysis method for solving the Schrodinger equation. *J Basic Appl Sci Res* 2012;2(6):6076–83.
- [12] Fariborzi Araghi MA, Naghshband S. On convergence of homotopy analysis method to solve the Schrodinger equation with a power law nonlinearity. *Int J Ind Math* 2013;5(4):367–74.
- [13] Ginibre J, Velo G. The cauchy problem in local spaces for the complex Ginzburg–Landau equation II. Contraction methods. *Commun Math Phys* 1997;187:45–79.
- [14] Gao H, Wang X. On the global existence and small dispersion limit for a class of complex Ginzburg–Landau equations. *Math Meth Appl Sci* 2009;32:1396–414.
- [15] He JH. Variational iteration method – some recent results and new interpretations. *J Comput Appl Math* 2007;207:3–17.
- [16] He JH. Variational iteration method a kind of non-linear analytical technique: some examples. *Int J Non-Linear Mech* 1999;34:699–708.
- [17] Irandoust-pakchin S, Kheiri H, Abdi-mazraeh S. Efficient computational algorithms for solving one class of fractional boundary value problems. *Comput Math Math Phys* 2013;53(7):920–32.
- [18] Jafarian A, Ghaderi P, Golmankhaneh AK, Baleanu D. Analytical treatment of system of Abel integral equations by homotopy analysis method. *Rom Rep Phys* 2014;66(3):603–11.
- [19] Jafarian A, Ghaderi P, Golmankhaneh AK, Baleanu D. Homotopy analysis method for solving coupled Ramani equations. *Rom J Phys* 2014;59(1–2):26–35.
- [20] Liao SJ. Beyond perturbation: introduction to the homotopy analysis method. Boca Raton: Chapman and Hall/CRC Press; 2003.
- [21] Liao SJ. Notes on the homotopy analysis method: some definitions and theorems. *Commun Nonlinear Sci Numer Simul* 2009;14:983–97.
- [22] Liao SJ. Homotopy analysis method in nonlinear differential equations. Springer; 2012.
- [23] Liao SJ. The proposed homotopy analysis technique for the solution of nonlinear problems. Ph.D. Thesis, Shanghai Jiao Tong University; 1992.
- [24] Marq P, Chate H, Conte R. Exact solutions of the one-dimensional quintic complex Ginzburg–Landau equation. *Physica D* 1994;73:305–17.
- [25] Ortega JB, Sardella E, Aguiar JA. Superconducting boundary conditions for mesoscopic circular samples. *Supercond Sci Technol* 2011;24(1):015001.
- [26] Ortega JB, Sardella E, Aguiar JA. Superconducting properties of a parallelepiped mesoscopic superconductor: a comparative study between the 2D and 3D Ginzburg–Landau models. *Phys Lett A* 2015;379(7):732–7.
- [27] Park J. Bifurcation and Stability of the generalized complex Ginzburg–Landau equation. *Pure Appl Anal* 2008;7(5):1237–53.
- [28] Rashidi MM, Domairry G, Dinarvand S. Approximate solutions for the Burger and regularized long wave equations by means of the homotopy analysis method. *Commun Nonlinear Sci Numer Simul* 2009;14:708–17.

- [29] Saarloos WV. The complex Ginzburg–Landau equation for beginners. P.O. Box 9506, 2300 RA, Leiden, The Netherlands: Instituut Lorentz, University of Leiden; 1994.
- [30] Sajid M, Hayat T, Asghar S. Comparison between the HAM and HPM solutions of thin film flows of non-Newtonian fluids on a moving belt. *Nonlinear Dyn* 2007;50(1–2):27–35.
- [31] Sherratt JA, Smith MJ, Rademacher JDM. Patterns of sources and sinks in the complex Ginzburg–Landau equation with zero linear dispersion. *SIAM J Appl Dyn Syst* 2010;9(3):883–918.
- [32] Siddheshwar PG. A series solution for the Ginzburg–Landau equation with a time-periodic coefficient. *Appl Math* 2010;3:542–54.
- [33] Wazwaz AM. Partial differential equations and solitary waves theory. Beijing and Springer-Verlag, Berlin Heidelberg: Higher Education Press; 2009.
- [34] Yang Y, Gao H. Continuous dependence on modelling for a complex Ginzburg–Landau equation with complex coefficients. *Math Meth Appl Sci* 2004;27:1567–78.



Soheila Naghshband was born in Tehran, in 1973. She received B.SC. degree in applied mathematics from Shahid Beheshti university and M.SC. degree in pure mathematics from Tehran university. She is the Ph.D. student of Islamic Azad university, Central Tehran branch. She is a faculty member of West Tehran branch, Islamic Azad university. Her research interests are numerical solution of nonlinear partial differential equations and related topics.



Mohammad Ali Fariborzi Araghi was born in Tehran, in 1967. He received B.SC. degree in mathematics from Tehran Teacher Training university, M.SC. degree in applied mathematics from Islamic Azad university, Karaj branch and Ph.D. degree in applied mathematics, numerical analysis field, from Islamic Azad university, Science and Research branch. He is the associate professor of Islamic Azad university, Central Tehran branch. His research interests are numerical solution of integral equations and partial differential equations and fuzzy mathematics.